

The More, The Merrier: The Who, What, and How of Group Therapy Across Patient Populations

2025 ASHA Convention Presentation by Jacalyn Savage

Reference List

- Archer, B., Tetnowski, J. C., Feer, S., & Schmadeke, E., & Christou-Franklin, E. (2019). Topic selection sequences in aphasia conversation groups. *Aphasiology*, 32(4), 394–416.
- Behrman, A., Cody, J., Elandary, S., Flom, P., & Chitnis, S. (2020). The effect of SPEAK OUT! and The LOUD Crowd on dysarthria due to Parkinson's disease. *American Journal of Speech-Language Pathology*, 29(3), 1448–1465.
- Bertcher, H. (1994). *Group participation: Techniques for leaders and members* (2nd ed.). Thousand Oaks, CA: Sage Publications.
- Bollinger, R., Musson, N., & Holland, A. (1993). A study of group communication intervention with chronically aphasic persons. *Aphasiology*, 7(4), 301–313.
- Bourque, K. S. (2020). Peer-mediated augmentative and alternative communication interventions for young children with autism spectrum disorder and limited to no spoken communication. *Perspectives of the ASHA Special Interest Groups*, 5(3), 602–610.
- Boyle, M., Akers, C. M., Cavanaugh, R., Hula, W. D., Swiderski, A. M., & Elman, E. J. (2023). Changes in discourse informativeness and efficiency following communication-based group treatment for chronic aphasia. *Aphasiology*, 37(3), 563–597.
- Connor, N. P., Orloff, D. I., & Uswatte, G. (2021). *Intensive social interaction for treatment of poststroke depression in subacute aphasia: The CONNECT Trial*. *Stroke*, 54(4), e189.
- Corbin, M. L. (1951). Group speech therapy for motor aphasia and dysarthria. *Journal of Speech and Hearing Disorders*, 16(1), 21–31.
- Elman, R. (Ed.). (1999a). *Group treatment of neurogenic communication disorders: The expert clinician's approach*. Boston, MA: Butterworth-Heinemann.
- Elman, R. (2000). *Working with groups: Neurogenic communication disorders and the managed care challenge* [Videotape]. Rockville, MD: American Speech-Language-Hearing Association.
- Elman, R. (Ed.). (2007a). *Group treatment of neurogenic communication disorders: The expert clinician's approach* (2nd ed.). San Diego, CA: Plural Publishing.
- Elman, R. (2007b). The importance of aphasia group treatment for rebuilding community and health. *Topics in Language Disorders*, 27(4), 300–308.
- Elman, R. (July, 2002). Aphasia groups: Healing through community. Keynote address presented at the International Aphasia Rehabilitation Conference, Brisbane, Australia.
- Ewing, S. (2007). Group process, group dynamics, and group techniques with neurogenic communication disorders. In R. Elman (Ed.), *Group treatment for neurogenic communication disorders: The expert clinician's approach* (2nd ed.). San Diego, CA: Plural Publishing.

Fama, J. M., Baron, C. R., Hatfield, B., & Turkeltaub, P. E. (2016). Effectiveness of group communication treatment in acute rehabilitation: A pilot study. *American Journal of Speech-Language Pathology*, 25(4), S798–S812.

Garrett, K., & Ellis, G. (1994). Group therapy for persons with long-term aphasia: Scaffolded language activities. *Aphasiology*, 8(5), 475–491.

Garrett, K., Staltari, C., & Moir, L. (2007). Contextual group communication therapy for persons with aphasia: A scaffolded discourse approach. In R. Elman (Ed.), *Group treatment for neurogenic communication disorders: The expert clinician's approach* (2nd ed.). San Diego, CA: Plural Publishing.

Gates, J. A., Kang, E., & Lerner, M. D. (2017). Efficacy of group social skills interventions for youth with autism spectrum disorder: A systematic review and meta-analysis. *Clinical Psychology Review*, 52, 164–181.

Hawley, L. A., & Newman, J. K. (2010). Group interactive structured treatment (GIST): A social competence intervention for individuals with brain injury. *Brain Injury*, 24(11), 1292–1297.

Hilari, K., & Northcott, S. (2006). Social support in people with chronic aphasia. *Aphasiology*, 20(1), 17–36.

Hoover, E. L., Caplan, D., Waters, G., & DeDe, G. (2020). Changes in connected speech and standardized tests following conversation treatment in chronic severe aphasia. *American Journal of Speech-Language Pathology*, 29(2), 808–823.

Hoover, E. L., DeDe, G., & Maas, E. (2021). *A randomized controlled trial of the effects of group conversation treatment on monologic discourse in aphasia*. *Journal of Speech, Language, and Hearing Research*, 64(12), 4861–4875.

Hoover, E. L., DeDe, G., & Maas, E. (2023). Multicenter randomized controlled trial of dyadic and group treatment for chronic aphasia. *Journal of Speech, Language, and Hearing Research*, 66(4), 1343–1364.

Hoover, E. L., Szabo, G., Kohen, F., Vitale, S., McCloskey, N., Maas, E., Kulkarni, V., & DeDe, G. (2025). *The benefits of conversation group treatment for individuals with chronic aphasia: Updated evidence from a multisite randomized controlled trial on measures of language and communication*. *American Journal of Speech-Language Pathology*, 34(3), 1203–1221.

Holland, A., & Ross, R. (1999). The power of aphasia groups. In R. Elman (Ed.), *Group treatment for neurogenic communication disorders: The expert clinician's approach* (pp. 115–117). Boston, MA: Butterworth-Heinemann.

Hung, S. L., Spencer, M. S., & Dronamraju, R. (2012). Selective mutism: Practice and intervention strategies for children. *Children and Schools*, 34(4), 222–230.

Hutchins, C., & Schmitt, M. B. (2024). Group size: An active ingredient of school-based language therapy. *Language, Speech, and Hearing Services in Schools*, 55(3), 781–802.

Kagan, A., & Simmons-Mackie, N. (2007). *The LIFE participation approach to aphasia: A clinician's guide*. Baltimore, MD: Brookes Publishing.

Kagan, A., Black, S., Duchan, J., Simmons-Mackie, N., & Square, P. (2001). Training volunteers as conversation partners using “Supported Conversation for Adults with Aphasia” (SCA): A controlled trial. *Journal of Speech, Language, and Hearing Research*, 44(3), 624–638.

- Kim, E., & Thompson, C. K. (2000). Patterns of discourse production in aphasia: A review of the literature. *Aphasiology*, 14(1), 43–64.
- Krajewska-Kulak, E., & Kozubski, W. (2020). A systematic review of active group-based dance, singing, music therapy and theatrical interventions for quality of life, functional communication, speech, motor function and cognitive status in people with Parkinson's disease. *Aging & Mental Health*, 24(12), 1964–1973.
- LaPointe, L. L., & Horner, J. (2010). Group therapy: Principles, methods, and strategies. In R. Chapey (Ed.), *Language intervention strategies in aphasia and related neurogenic communication disorders* (5th ed., pp. 609–626). Baltimore, MD: Lippincott Williams & Wilkins.
- MacWhinney, B. (2000). *The CHILDES project: Tools for analyzing talk* (3rd ed.). Mahwah, NJ: Lawrence Erlbaum Associates.
- Maier, L., & Swain, A. (2012). Social connectedness and quality of life in aphasia: A systematic review. *Aphasiology*, 26(11), 1316–1335.
- Maul, R., & Hall, T. (2021). Peer-mediated group interventions for children with communication disorders: Evidence and recommendations. *Communication Disorders Quarterly*, 42(3), 148–158.
- McVay, J. C., & Mooney, R. A. (2006). Aphasia groups: Making therapy meaningful. *Topics in Stroke Rehabilitation*, 13(2), 1–12.
- Meteyard, L., & Holmes, N. (2020). Best practices in group interventions for individuals with neurogenic communication disorders. *Perspectives of the ASHA Special Interest Groups*, 5(6), 1387–1399.
- Murray, L. L. (2012). The rehabilitation of aphasia: Current approaches and future directions. *Language and Linguistics Compass*, 6(12), 739–758.
- Northcott, S., Marshall, J., & Hilari, K. (2016). Psychosocial outcomes in people with chronic aphasia attending community-based group interventions. *Aphasiology*, 30(12), 1460–1479.
- Northcott, S., & Hilari, K. (2011). The impact of stroke on social participation and quality of life: Integrating the evidence. *International Journal of Speech-Language Pathology*, 13(4), 309–315.
- Parr, S. (2007). Living with severe aphasia: Tracking social exclusion. *Aphasiology*, 21(1), 98–123.
- Pulvermüller, F., & Berthier, M. L. (2008). Aphasia therapy on a neuroscience basis. *Aphasiology*, 22(6), 563–599.
- Ramsberger, G., & Rende, R. (2002). Effects of structured group conversation therapy on verbal communication in individuals with chronic aphasia. *Aphasiology*, 16(10–11), 1007–1023.
- Rinaldi, M., & Basil, E. (2023). Technology-supported group interventions for speech and language disorders. *Frontiers in Rehabilitation Sciences*, 4, 1145829.
- Simmons-Mackie, N., Elman, R., Holland, A., & Damico, J. (2007). Management of discourse in group therapy for aphasia. *Topics in Language Disorders*, 27(1), 4–22.
- Searl, J., Wilson, K., Haring, K., Dietsch, A., Lyons, K., & Pahwa, R. (2011). Feasibility of group voice therapy for individuals with Parkinson's disease. *Journal of Communication Disorders*, 44(6), 719–732.

- Spencer, K., et al. (2001). Evidence-based practice guidelines for dysarthria. American Neurogenic Communication Disorders Society (ANCDS).
- Staltari, C., Garrett, K., & Moir, L. (2007). Contextual Group Communication Therapy for persons with aphasia: A scaffolded discourse approach. In R. Elman (Ed.), *Group treatment for neurogenic communication disorders: The expert clinician's approach* (2nd ed.). San Diego, CA: Plural Publishing.
- Taylor, N., & Francis, D. (2020). Communication-oriented group therapy for dysarthria: A randomized controlled trial. *International Journal of Language & Communication Disorders*, 55(4), 512–528.
- Thiemann-Bourque, K., & Brady, N. (2018). Incorporating a peer-mediated approach into speech-generating device (SGD) intervention in preschoolers with autism. *American Journal of Speech-Language Pathology*, 27(3), 899–911.
- Thiemann-Bourque, K., Brady, N., & Fleming, K. (2020). Peer-mediated augmentative and alternative communication intervention: Children with ASD communicate more using SGDs after peer training. *Perspectives of the ASHA Special Interest Groups*, 5(3), 602–610.
- Togher, L., et al. (2023). INCOG 2.0 guidelines for cognitive rehabilitation: Social communication. *The INCOG 2.0 recommendations reflect group treatments to improve social communication*. (PubMed)
- Togher, L., et al. (2013). Non-randomized trial of communication partner training + group + individual for TBI: *J Rehabil Med*, 45, 637–645.
- Tuckman, B. W. (1965). Developmental sequence in small groups. *Psychological Bulletin*, 63(6), 384–399.
- Vickers, C. P. (2010). Social networks after the onset of aphasia: The impact of aphasia group attendance. *Aphasiology*, 24(6–8), 902–913.
- Watanabe, M., Cartwright, J., & Pierce, J. E. (2024). Positive effects of speech and language therapy group interventions in primary progressive aphasia: A systematic review. *International Journal of Language & Communication Disorders*, 59(5), 1832–1849.